

Southern Illinois University Carbondale OpenSIUC

Miscellaneous (presentations, translations,
interviews, etc)

Department of Mathematics

8-1998

On the Dynamics of Stochastic Differential Systems (The Seventh Vilnius Conference on Probability Theory and Mathematical Statistics)

Salah-Eldin A. Mohammed

Southern Illinois University Carbondale, salah@sfde.math.siu.edu

Follow this and additional works at: http://opensiuc.lib.siu.edu/math_misc

 Part of the [Mathematics Commons](#)

The Seventh Vilnius Conference on Probability Theory and Mathematical Statistics; August 12-18, 1998; Vilnius, Lithuania

Recommended Citation

Mohammed, Salah-Eldin A., "On the Dynamics of Stochastic Differential Systems (The Seventh Vilnius Conference on Probability Theory and Mathematical Statistics)" (1998). *Miscellaneous (presentations, translations, interviews, etc)*. Paper 25.
http://opensiuc.lib.siu.edu/math_misc/25

This Article is brought to you for free and open access by the Department of Mathematics at OpenSIUC. It has been accepted for inclusion in Miscellaneous (presentations, translations, interviews, etc) by an authorized administrator of OpenSIUC. For more information, please contact opensiuc@lib.siu.edu.

**ON THE DYNAMICS OF STOCHASTIC
DIFFERENTIAL SYSTEMS**

Vilnius : August 13, 1998

Salah-Eldin A. Mohammed

Southern Illinois University
Carbondale, IL 62901-4408 USA

Web site: <http://salah.math.siu.edu>

Outline

- Formulate a *Local Stable Manifold Theorem* for stochastic differential equations with and without memory (SFDE's and SODE's).
- Spatial Kunita-type semimartingales noise, with stationary ergodic increments.
- Start with the existence of a stochastic semiflow for SDE.
- Concept of a hyperbolic stationary trajectory. For Stratonovich SODE, stationary trajectory is a solution of the forward /backward anticipating equation for all time.
- Existence of a stationary random family of asymptotically invariant stable and unstable manifolds within a stationary neighborhood of the hyperbolic stationary solution.
- For Stratonovich SODE, stable and unstable manifolds are dynamically characterized using forward and backward solutions of anticipating versions of the equation.

- Proofs based on Ruelle-Oseledec (non-linear) multiplicative ergodic theory and anticipating stochastic calculus.

Formulation of the Theorem

I. SODE Case:

Stratonovich SODE

$$dx(t) = h(x(t)) dt + \sum_{i=1}^m g_i(x(t)) \circ dW_i(t), \quad (I)$$

on \mathbf{R}^d driven by m -dimensional Brownian motion $W := (W_1, \dots, W_m)$.

$(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbf{R}}, P) :=$ canonical filtered Wiener space.

$\Omega :=$ space of all continuous paths $\omega : \mathbf{R} \rightarrow \mathbf{R}^m$, $\omega(0) = 0$, in Euclidean space \mathbf{R}^m , with compact open topology;

$\mathcal{F} :=$ Borel σ -field of Ω ;

$\mathcal{F}_t :=$ sub- σ -field of \mathcal{F} generated by the evaluations $\omega \rightarrow \omega(u)$, $u \leq t$, $t \in \mathbf{R}$.

$P :=$ Wiener measure on Ω .

$h, g_i : \mathbf{R}^d \rightarrow \mathbf{R}^d$, $1 \leq i \leq m$, vector fields on \mathbf{R}^d . For some $k \geq 1, \delta \in (0, 1)$, h is $C_b^{k, \delta}$, viz. h has all derivatives $D^j h$, $1 \leq j \leq k$, continuous and globally bounded, $D^k h$ Hölder continuous with exponent δ . g_i , $1 \leq i \leq m$, globally bounded and $C_b^{k+1, \delta}$.

$\theta : \mathbf{R} \times \Omega \rightarrow \Omega$ is the (ergodic) Brownian shift

$$\theta(t, \omega)(s) := \omega(t + s) - \omega(t), \quad t, s \in \mathbf{R}, \omega \in \Omega.$$


Let $\phi : \mathbf{R} \times \mathbf{R}^d \times \Omega \rightarrow \mathbf{R}^d$ be the stochastic flow of (I) ($\phi(t, \cdot, \omega) = [\phi(-t, \cdot, \theta(t, \omega))]^{-1}, t < 0$). Then ϕ is a perfect $C^{k, \epsilon}$ cocycle:

$$\phi(t + s, \cdot, \omega) = \phi(t, \cdot, \theta(s, \omega)) \circ \phi(s, \cdot, \omega),$$

for all $s, t \in \mathbf{R}$ and all $\omega \in \Omega, \epsilon \in (0, \delta)$ ([I-W], [Ku], [A-S]).

Figure illustrates the cocycle property. Vertical solid lines represent random fibers consisting of copies of \mathbf{R}^d (or a Banach space of paths in \mathbf{R}^d .) (ϕ, θ) is a “random vector-bundle morphism” over the “base” probability space Ω .

The Cocycle



Definition

SODE (I) has a *stationary point* if there exists an \mathcal{F} -measurable random variable $Y : \Omega \rightarrow \mathbf{R}^d$ such that

$$\phi(t, Y(\omega), \omega) = Y(\theta(t, \omega)) \quad (1)$$

for all $t \in \mathbf{R}$ and every $\omega \in \Omega$. Denote stationary trajectory (1) by $\phi(t, Y) = Y(\theta(t))$.

Examples of Stationary Points

1. Fixed points:

$$d\phi(t) = h(\phi(t)) dt + \sum_{i=1}^m g_i(\phi(t)) \circ dW_i(t)$$

$$h(x_0) = g_i(x_0) = 0, \quad 1 \leq i \leq m$$

Take $Y(\omega) = x_0$ for all $\omega \in \Omega$.

2. Linear affine case $d = 1$:

$$d\phi(t) = \lambda\phi(t) dt + dW(t)$$

$\lambda > 0$ fixed, $W(t) \in \mathbf{R}$. Take

$$Y(\omega) := - \int_0^\infty e^{-\lambda u} dW(u),$$

$$\theta(t, \omega)(s) = \omega(t + s) - \omega(t).$$

Check that $\phi(t, Y(\omega), \omega) = Y(\theta(t, \omega))$, using integration by parts and variation of parameters.

3. Affine linear SODE in $d = 2$:

$$d\phi(t) = A\phi(t) dt + GdW(t)$$

with A a fixed hyperbolic 2×2 -diagonal matrix; G a constant matrix.

4. Generate a large class of stationary points as follows: Let ρ be an invariant probability measure for the one-point motion in \mathbf{R}^d . Then ρ gives rise to a stationary point by suitably enlarging the underlying probability space: If $P_t : C_b(\mathbf{R}^d, \mathbf{R}) \rightarrow C_b(\mathbf{R}^d, \mathbf{R}), t \geq 0$, is the Markov semigroup associated with the SODE, then

$$\int_{\mathbf{R}^d} (P_t f)(x) d\rho(x) = \int_{\mathbf{R}^d} f(x) d\rho(x), \quad t \geq 0$$

where

$$(P_t f)(x) := E[f(\phi(t, x, \cdot))], \quad t \geq 0, \quad x \in \mathbf{R}^d,$$

for all $f \in C_b(\mathbf{R}^d, \mathbf{R})$. Define

$$\begin{aligned}\tilde{\Omega} &:= \Omega \times \mathbf{R}^d, & \tilde{\mathcal{F}} &:= \mathcal{F} \otimes \mathcal{B}(\mathbf{R}^d), & \tilde{P} &:= P \otimes \rho, & \tilde{\omega} &:= (\omega, x) \in \tilde{\Omega}, \\ \tilde{\theta}(t, \tilde{\omega}) &:= (\theta(t, \omega), \phi(t, x, \omega)), & t &\in \mathbf{R}^+, \omega \in \Omega, x \in \mathbf{R}^d \\ \tilde{\phi}(t, x', \tilde{\omega}) &:= \phi(t, x', \omega), & t &\in \mathbf{R}^+, x' \in \mathbf{R}^d, \tilde{\omega} \in \tilde{\Omega} \\ \tilde{Y}(\tilde{\omega}) &:= x, & \tilde{\omega} &= (\omega, x) \in \tilde{\Omega}.\end{aligned}$$

The group $\tilde{\theta}(t, \cdot) : \tilde{\Omega} \rightarrow \tilde{\Omega}, t \in \mathbf{R}^+$, is \tilde{P} -preserving (and ergodic) (Carverhill [C]). $(\tilde{\phi}(t, \cdot, \tilde{\omega}), \tilde{\theta}(t, \tilde{\omega}))$ is a perfect cocycle on \mathbf{R}^d ; and $\tilde{Y} : \tilde{\Omega} \rightarrow \mathbf{R}^d$ satisfies

$$\tilde{\phi}(t, \tilde{Y}(\tilde{\omega}), \tilde{\omega}) = \tilde{Y}(\tilde{\theta}(t, \tilde{\omega}))$$

for all $t \in \mathbf{R}^+, \tilde{\omega} \in \tilde{\Omega}$. Hence \tilde{Y} is a stationary point for the cocycle $(\tilde{\phi}, \tilde{\theta})$, and $\rho = \tilde{P} \circ \tilde{Y}^{-1}$.

Conversely, let $Y : \Omega \rightarrow \mathbf{R}^d$ be a stationary random point satisfying the identity (1) and independent of the Brownian motion $W(t), t \geq 0$. Then $\rho := P \circ Y^{-1}$ is an invariant measure for the one-point motion.

Let $\phi(t, Y)$ be a stationary solution of (I). Cocycle property of ϕ implies that the linearization

$$(D_2\phi(t, Y(\omega), \omega), \theta(t, \omega))$$

along the stationary solution is also a $d \times d$ -matrix-valued cocycle. Using Kolmogorov's theorem, the random variables

$$\sup_{x \in \mathbf{R}^d} \frac{|D_2\phi(t, x)|}{(1 + |x|^\gamma)}, \quad \gamma > 0,$$

have moments of all orders. If $E \log^+ |Y| < \infty$, then $E \log^+ |D_2\phi(1, Y)| < \infty$. Apply Oseledec's Theorem to get a *non-random* finite Lyapunov spectrum:

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log |D_2\phi(n, Y(\omega), \omega)(v(\omega))|, \quad v \in L^0(\Omega, \mathbf{R}^d).$$

Spectrum takes finitely many values $\{\lambda_i\}_{i=1}^p$ with non-random multiplicities q_i , $1 \leq i \leq p$, and $\sum_{i=1}^p q_i = d$ ([Ru.1], Theorem I.6).

Definition

A stationary point Y is *hyperbolic* if $E \log^+ |Y(\cdot)| < \infty$, and if $(D_2\phi(n, Y(\omega), \omega), \theta(n, \omega))$ has a non-vanishing Lyapunov spectrum

$$\{\lambda_p < \cdots < \lambda_{i_0+1} < \lambda_{i_0} < 0 < \lambda_{i_0-1} < \cdots < \lambda_2 < \lambda_1\}$$

i.e. $\lambda_i \neq 0$ for all $1 \leq i \leq p$.

Define $\lambda_{i_0} := \max\{\lambda_i : \lambda_i < 0\}$ if at least one $\lambda_i < 0$. If all $\lambda_i > 0$, set $\lambda_{i_0} = -\infty$. (This implies that $\lambda_{i_0-1} := \min\{\lambda_i :$

$\lambda_i > 0\}$, if at least one $\lambda_i > 0$; in case all λ_i are negative, set $\lambda_{i_0-1} = \infty$.)

Let $\rho \in \mathbf{R}^+$, $x \in \mathbf{R}^d$.

$B(x, \rho) :=$ open ball in \mathbf{R}^d , center x and radius ρ ;

$\bar{B}(x, \rho) :=$ corresponding closed ball;

$\mathcal{C}(\mathbf{R}^d) :=$ the class of all non-empty compact subsets of \mathbf{R}^d with Hausdorff metric d^* :

$$d^*(A_1, A_2) := \sup\{d(x, A_1) : x \in A_2\} \vee \sup\{d(y, A_2) : y \in A_1\}$$

where $A_1, A_2 \in \mathcal{C}(\mathbf{R}^d)$;

$d(x, A_i) := \inf\{|x - y| : y \in A_i\}$, $x \in \mathbf{R}^d$, $i = 1, 2$;

$\mathcal{B}(\mathcal{C}(\mathbf{R}^d)) :=$ Borel σ -algebra on $\mathcal{C}(\mathbf{R}^d)$ with respect to the metric d^* .

Theorem 1 (Stable Manifold Theorem-SODE) ([M-S], 1997)

Assume that the coefficients of SODE (I) satisfy the given hypotheses. Suppose $\phi(t, Y)$ is a hyperbolic stationary trajectory of (I) with $E \log^+ |Y| < \infty$.

Fix $\epsilon_1 \in (0, -\lambda_{i_0})$ and $\epsilon_2 \in (0, \lambda_{i_0-1})$. Then there exist

- (i) a sure event $\Omega^* \in \mathcal{F}$ with $\theta(t, \cdot)(\Omega^*) = \Omega^*$ for all $t \in \mathbf{R}$,
- (ii) \mathcal{F} -measurable random variables $\rho_i, \beta_i : \Omega^* \rightarrow (0, 1)$, $\beta_i > \rho_i > 0$, $i = 1, 2$, such that for each $\omega \in \Omega^*$, the following is true:

There are $C^{k, \epsilon}$ ($\epsilon \in (0, \delta)$) submanifolds $\tilde{\mathcal{S}}(\omega), \tilde{\mathcal{U}}(\omega)$ of $\bar{B}(Y(\omega), \rho_1(\omega))$ and $\bar{B}(Y(\omega), \rho_2(\omega))$ (resp.) with the following properties:

- (a) $\tilde{\mathcal{S}}(\omega)$ is the set of all $x \in \bar{B}(Y(\omega), \rho_1(\omega))$ such that

$$|\phi(n, x, \omega) - Y(\theta(n, \omega))| \leq \beta_1(\omega) e^{(\lambda_{i_0} + \epsilon_1)n}$$

for all integers $n \geq 0$. Furthermore,

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log |\phi(t, x, \omega) - Y(\theta(t, \omega))| \leq \lambda_{i_0} \quad (2)$$

for all $x \in \tilde{\mathcal{S}}(\omega)$. Each stable subspace $\mathcal{S}(\omega)$ of the linearized flow $D_2\phi$ is tangent at $Y(\omega)$ to the submanifold $\tilde{\mathcal{S}}(\omega)$, viz. $T_{Y(\omega)}\tilde{\mathcal{S}}(\omega) = \mathcal{S}(\omega)$. In particular, $\dim \tilde{\mathcal{S}}(\omega) = \dim \mathcal{S}(\omega)$ and is non-random.

$$(b) \limsup_{t \rightarrow \infty} \frac{1}{t} \log \left[\sup_{\substack{x_1 \neq x_2 \\ x_1, x_2 \in \tilde{\mathcal{S}}(\omega)}} \left\{ \frac{|\phi(t, x_1, \omega) - \phi(t, x_2, \omega)|}{|x_1 - x_2|} \right\} \right] \leq \lambda_{i_0}.$$

(c) (Cocycle-invariance of the stable manifolds):

There exists $\tau_1(\omega) \geq 0$ such that

$$\phi(t, \cdot, \omega)(\tilde{\mathcal{S}}(\omega)) \subseteq \tilde{\mathcal{S}}(\theta(t, \omega)), \quad t \geq \tau_1(\omega). \quad (3)$$

Also

$$D_2\phi(t, Y(\omega), \omega)(\mathcal{S}(\omega)) = \mathcal{S}(\theta(t, \omega)), \quad t \geq 0. \quad (4)$$

(d) $\tilde{\mathcal{U}}(\omega)$ is the set of all $x \in \bar{B}(Y(\omega), \rho_2(\omega))$ with the property that

$$|\phi(-n, x, \omega) - Y(\theta(-n, \omega))| \leq \beta_2(\omega) e^{(-\lambda_{i_0-1} + \epsilon_2)n} \quad (5)$$

for all integers $n \geq 0$. Also

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log |\phi(-t, x, \omega) - Y(\theta(-t, \omega))| \leq -\lambda_{i_0-1}. \quad (6)$$

for all $x \in \tilde{\mathcal{U}}(\omega)$. Furthermore, the unstable subspace $\mathcal{U}(\omega)$ of $D_2\phi$ is the tangent space to $\tilde{\mathcal{U}}(\omega)$ at $Y(\omega)$, viz. $T_{Y(\omega)}\tilde{\mathcal{U}}(\omega) = \mathcal{U}(\omega)$. In particular, $\dim \tilde{\mathcal{U}}(\omega) = \dim \mathcal{U}(\omega)$ and is non-random.

$$(e) \limsup_{t \rightarrow \infty} \frac{1}{t} \log \left[\sup_{\substack{x_1 \neq x_2 \\ x_1, x_2 \in \tilde{\mathcal{U}}(\omega)}} \left\{ \frac{|\phi(-t, x_1, \omega) - \phi(-t, x_2, \omega)|}{|x_1 - x_2|} \right\} \right] \leq -\lambda_{i_0-1}.$$

(f) (Cocycle-invariance of the unstable manifolds):

There exists $\tau_2(\omega) \geq 0$ such that

$$\phi(-t, \cdot, \omega)(\tilde{\mathcal{U}}(\omega)) \subseteq \tilde{\mathcal{U}}(\theta(-t, \omega)), \quad t \geq \tau_2(\omega). \quad (7)$$

Also

$$D_2\phi(-t, Y(\omega), \omega)(\mathcal{U}(\omega)) = \mathcal{U}(\theta(-t, \omega)), \quad t \geq 0. \quad (8)$$

(g) The submanifolds $\tilde{\mathcal{U}}(\omega)$ and $\tilde{\mathcal{S}}(\omega)$ are transversal, viz.


$$\mathbf{R}^d = T_{Y(\omega)}\tilde{\mathcal{U}}(\omega) \oplus T_{Y(\omega)}\tilde{\mathcal{S}}(\omega). \quad (9)$$

(h) The mappings

$$\begin{aligned} \Omega &\rightarrow \mathcal{C}(\mathbf{R}^d), & \Omega &\rightarrow \mathcal{C}(\mathbf{R}^d), \\ \omega &\mapsto \tilde{\mathcal{S}}(\omega) & \omega &\mapsto \tilde{\mathcal{U}}(\omega) \end{aligned}$$


are $(\mathcal{F}, \mathcal{B}(\mathcal{C}(\mathbf{R}^d)))$ -measurable.

Assume, further, that $h, g_i, 1 \leq i \leq m$, are C_b^∞ . Then the local stable and unstable manifolds $\tilde{\mathcal{S}}(\omega), \tilde{\mathcal{U}}(\omega)$ are C^∞ .



$$t > \tau_1(\omega)$$

A picture is worth a 1000 words!



$$t > \tau_2(\omega)$$

II. SFDE Case:

“Regular” Itô SFDE with finite memory:

$$\left. \begin{aligned} dx(t) &= H(x(t), x_t) dt + \sum_{i=1}^m G_i(x(t), g(x_t)) dW_i(t), \\ (x(0), x_0) &= (v, \eta) \in M_2 := \mathbf{R}^d \times L^2([-r, 0], \mathbf{R}^d) \end{aligned} \right\} \quad (I')$$

Solution segment $x_t(s) := x(t + s)$, $t \geq 0, s \in [-r, 0]$.

Smooth memory: $g : L^2([-r, 0], \mathbf{R}^d) \rightarrow \mathbf{R}^p$ is $C^{k,\delta}$ ($k \geq 1, \delta \in (0, 1]$) with all Fréchet derivatives $D^j g, 1 \leq j \leq k$, globally bounded; $t \rightarrow g_i(x_t)$ locally of B.V., with

$$L^2([-r, T], \mathbf{R}^d) \ni x \mapsto \{t \mapsto \frac{dg(x_t)}{dt}\} \in L^2([0, T], \mathbf{R}^m)$$

globally bounded, globally Lipschitz and of class $C^{k,\delta}$.

m -dimensional Brownian motion $W := (W_1, \dots, W_m)$

Ergodic Brownian shift θ as before.

State space M_2 , Hilbert with usual norm.

Smoothness Hypotheses:

$H : M_2 \rightarrow \mathbf{R}^d$ of class $C_b^{k,\delta}$, viz. all Fréchet derivatives $D^j H, 1 \leq j \leq k$, continuous and globally bounded, $D^k H$ Hölder continuous with exponent δ on bounded sets in M_2 .

$G_i : \mathbf{R}^d \times \mathbf{R}^p \rightarrow \mathbf{R}^d, 1 \leq i \leq m$, of class $C^{k+1,\delta}$.

Then (I') has a stochastic semiflow $X : \mathbf{R}^+ \times M_2 \times \Omega \rightarrow M_2$ with $X(t, (v, \eta), \cdot) = (x(t), x_t)$. X is of class $C^{k,\epsilon}$ for any $\epsilon \in (0, \delta)$, takes bounded sets into relatively compact sets in M_2 . (X, θ) is a perfect cocycle on M_2 ([M-S]). Define *hyperbolic stationary point* $Y : \Omega \rightarrow M_2$ as in SODE case (replace ϕ by X ; apply [Mo.1] to linearized cocycle).

Theorem 1' (Stable Manifold Theorem-SFDE) ([M-S], 1998)

Assume above smoothness Hypotheses on H, G_i, g . Let Y be a hyperbolic stationary point of (I') such that $E(\|Y(\cdot)\|^{\epsilon_0}) < \infty$ for some $\epsilon_0 > 0$

Suppose $(D_2X(t, Y(\omega), \omega), \theta(t, \omega), t \geq 0)$ has a Lyapunov spectrum $\{\dots < \lambda_{i+1} < \lambda_i < \dots < \lambda_2 < \lambda_1\}$. Define λ_{i_0} to be the largest negative Lyapunov exponent (as before).

Fix $\epsilon_1 \in (0, -\lambda_{i_0})$ and $\epsilon_2 \in (0, \lambda_{i_0-1})$. Then there exist

- (i) a sure event $\Omega^* \in \mathcal{F}$ with $\theta(t, \cdot)(\Omega^*) = \Omega^*$ for all $t \in \mathbf{R}$,
- (ii) \mathcal{F} -measurable random variables $\rho_i, \beta_i : \Omega^* \rightarrow (0, 1)$, $\beta_i > \rho_i > 0$, $i = 1, 2$, such that for each $\omega \in \Omega^*$, the following is true:

There are $C^{k,\epsilon}$ ($\epsilon \in (0, \delta)$) submanifolds $\tilde{S}(\omega), \tilde{U}(\omega)$ of $\bar{B}(Y(\omega), \rho_1(\omega))$ and $\bar{B}(Y(\omega), \rho_2(\omega))$ (resp.) with the following properties:

(a) $\tilde{\mathcal{S}}(\omega)$ is the set of all $(v, \eta) \in \bar{B}(Y(\omega), \rho_1(\omega))$ such that

$$\|X(n, (v, \eta), \omega) - Y(\theta(n, \omega))\| \leq \beta_1(\omega) e^{(\lambda_{i_0} + \epsilon_1)n}$$

for all integers $n \geq 0$. Furthermore,

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log \|X(t, (v, \eta), \omega) - Y(\theta(t, \omega))\| \leq \lambda_{i_0}$$

for all $(v, \eta) \in \tilde{\mathcal{S}}(\omega)$. Each stable subspace $\mathcal{S}(\omega)$ of the linearized semiflow D_2X is tangent at $Y(\omega)$ to the submanifold $\tilde{\mathcal{S}}(\omega)$, viz. $T_{Y(\omega)}\tilde{\mathcal{S}}(\omega) = \mathcal{S}(\omega)$. In particular, $\text{codim } \tilde{\mathcal{S}}(\omega) = \text{codim } \mathcal{S}(\omega)$, is fixed and finite.

$$(b) \limsup_{t \rightarrow \infty} \frac{1}{t} \log \left[\sup \left\{ \frac{\|X(t, (v_1, \eta_1), \omega) - X(t, (v_2, \eta_2), \omega)\|}{\|(v_1, \eta_1) - (v_2, \eta_2)\|} : (v_1, \eta_1) \neq (v_2, \eta_2), (v_1, \eta_1), (v_2, \eta_2) \in \tilde{\mathcal{S}}(\omega) \right\} \right] \leq \lambda_{i_0}.$$

(c) (Cocycle-invariance of the stable manifolds):

There exists $\tau_1(\omega) \geq 0$ such that

$$X(t, \cdot, \omega)(\tilde{\mathcal{S}}(\omega)) \subseteq \tilde{\mathcal{S}}(\theta(t, \omega)), \quad t \geq \tau_1(\omega).$$

Also

$$D_2X(t, Y(\omega), \omega)(\mathcal{S}(\omega)) = \mathcal{S}(\theta(t, \omega)), \quad t \geq 0.$$

(d) $\tilde{\mathcal{U}}(\omega)$ is the set of all $(v, \eta) \in \bar{B}(Y(\omega), \rho_2(\omega))$ with the property that there is a unique “history” process $y(\cdot, \omega) : \mathbf{Z}^- \rightarrow M_2$

such that $y(0, \omega) = (v, \eta)$ and for each integer $n \geq 1$, one has $X(r, y(-nr, \omega), \theta(-nr, \omega)) = y(-(n-1)r, \omega)$ and

$$\|y(-nr, \omega) - Y(\theta(-nr, \omega))\|_{M_2} \leq \beta_2(\omega) e^{-(\lambda_{i_0-1} - \epsilon_2)nr}.$$

Furthermore, for each $(v, \eta) \in \tilde{\mathcal{U}}(\omega)$, there is a unique continuous-time “history” process also denoted by $y(\cdot, \omega) : (-\infty, 0] \rightarrow M_2$ such that $y(0, \omega) = (v, \eta)$, $X(t, y(s, \omega), \theta(s, \omega)) = y(t+s, \omega)$ for all $s \leq 0, 0 \leq t \leq -s$, and

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log \|y(-t, \omega) - Y(\theta(-t, \omega))\| \leq -\lambda_{i_0-1}.$$

Each unstable subspace $\mathcal{U}(\omega)$ of the linearized semiflow D_2X is tangent at $Y(\omega)$ to $\tilde{\mathcal{U}}(\omega)$, viz. $T_{Y(\omega)}\tilde{\mathcal{U}}(\omega) = \mathcal{U}(\omega)$. In particular, $\dim \tilde{\mathcal{U}}(\omega)$ is finite and non-random.

(e) Let $y(\cdot, (v_i, \eta_i), \omega), i = 1, 2$, be the history processes associated with $(v_i, \eta_i) = y(0, (v_i, \eta_i), \omega) \in \tilde{\mathcal{U}}(\omega), i = 1, 2$. Then

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log \left[\sup \left\{ \frac{\|y(-t, (v_1, \eta_1), \omega) - y(-t, (v_2, \eta_2), \omega)\|}{\|(v_1, \eta_1) - (v_2, \eta_2)\|} : \right. \right. \\ \left. \left. (v_1, \eta_1) \neq (v_2, \eta_2), (v_i, \eta_i) \in \tilde{\mathcal{U}}(\omega), i = 1, 2 \right\} \right] \leq -\lambda_{i_0-1}.$$

(f) (Cocycle-invariance of the unstable manifolds):

There exists $\tau_2(\omega) \geq 0$ such that

$$\tilde{\mathcal{U}}(\omega) \subseteq X(t, \cdot, \theta(-t, \omega))(\tilde{\mathcal{U}}(\theta(-t, \omega)))$$

for all $t \geq \tau_2(\omega)$. Also

$$D_2X(t, \cdot, \theta(-t, \omega))(\mathcal{U}(\theta(-t, \omega))) = \mathcal{U}(\omega), \quad t \geq 0;$$

and the restriction

$$D_2X(t, \cdot, \theta(-t, \omega))|_{\mathcal{U}(\theta(-t, \omega))} : \mathcal{U}(\theta(-t, \omega)) \rightarrow \mathcal{U}(\omega), \quad t \geq 0,$$

is a linear homeomorphism onto.

(g) The submanifolds $\tilde{\mathcal{U}}(\omega)$ and $\tilde{\mathcal{S}}(\omega)$ are transversal, viz.

$$M_2 = T_{Y(\omega)}\tilde{\mathcal{U}}(\omega) \oplus T_{Y(\omega)}\tilde{\mathcal{S}}(\omega).$$

(h) The mappings

$$\begin{aligned} \Omega &\rightarrow \mathcal{C}(M_2), & \Omega &\rightarrow \mathcal{C}(M_2), \\ \omega &\mapsto \tilde{\mathcal{S}}(\omega) & \omega &\mapsto \tilde{\mathcal{U}}(\omega) \end{aligned}$$

are $(\mathcal{F}, \mathcal{B}(\mathcal{C}(M_2)))$ -measurable.

Assume, in addition, that the smoothness hypotheses hold for every $k \geq 1$ and $\delta \in (0, 1]$. Then the local stable and unstable manifolds $\tilde{\mathcal{S}}(\omega)$, $\tilde{\mathcal{U}}(\omega)$ are C^∞ .

Sketch of Proof-SODE Case

Broad Outline:

- Linearize along stationary solution and use substitution formula:

$$\left. \begin{aligned} d\phi(t, Y) &= h(\phi(t, Y)) dt + \sum_{i=1}^m g_i(\phi(t, Y)) \circ dW_i(t), & t > 0 \\ \phi(0, Y) &= Y. \end{aligned} \right\} \quad (II)$$

([N-P]).

$$\left. \begin{aligned} dD_2\phi(t, Y) &= Dh(\phi(t, Y))D_2\phi(t, Y) dt \\ &\quad + \sum_{i=1}^m Dg_i(\phi(t, Y))D_2\phi(t, Y) \circ dW_i(t), & t > 0 \\ D_2\phi(0, Y) &= I. \end{aligned} \right\} \quad (III)$$

D_2, D denotes spatial (Fréchet) derivatives.

$$\left. \begin{aligned} d\phi(t, Y) &= -h(\phi(t, Y)) dt - \sum_{i=1}^m g_i(\phi(t, Y)) \circ \hat{d}W_i(t), & t < 0 \\ \phi(0, Y) &= Y. \end{aligned} \right\} \quad (II^-)$$

$$\left. \begin{aligned} dD_2\phi(t, Y) &= -Dh(\phi(t, Y))D_2\phi(t, Y) dt \\ &\quad - \sum_{i=1}^m Dg_i(\phi(t, Y))D_2\phi(t, Y) \circ \hat{d}W_i(t), & t < 0 \\ D_2\phi(0, Y) &= I. \end{aligned} \right\} \quad (III^-)$$

Above SODE's (II)-(III)⁻ give dynamic characterizations of stable/unstable manifolds.

- “Perfection” of ergodic theorem and Kingman’s sub-additive ergodic theorem under suitable integrability hypotheses.
- Apply the Oseledec theorem to the linearized system. Get a fixed Lyapunov spectrum. Hyperbolicity is well-defined.
- Continuous-time integrability estimates on the non-linear cocycle in a neighborhood of the stationary trajectory. Uses sharp global spatial estimates on the stochastic flow via Kolmogorov’s theorem; viz. the following quantities have q -th moments for all $q \geq 1$:

$$\begin{aligned} & \sup_{\substack{0 \leq s, t \leq T, \\ x \in \mathbf{R}^d}} \frac{|\phi_{s,t}(x, \omega)|}{[1 + |x|(\log^+ |x|)^\gamma]}, \quad \sup_{\substack{0 \leq s, t \leq T, \\ x \in \mathbf{R}^d}} \frac{|D_x^\alpha \phi_{s,t}(x, \omega)|}{(1 + |x|^\gamma)}, \\ & \sup_{x \in \mathbf{R}^d} \sup_{\substack{0 \leq s, t \leq T, \\ 0 < |x' - x| \leq \rho}} \frac{|D_x^\alpha \phi_{s,t}(x, \omega) - D_x^\alpha \phi_{s,t}(x', \omega)|}{|x - x'|^\epsilon (1 + |x|)^\gamma}, \\ & \epsilon \in (0, \delta), \gamma, \rho, T > 0, 1 \leq |\alpha| \leq k \end{aligned}$$

- Use Ruelle’s discrete non-linear ergodic theorem on the auxiliary perfect cocycle

$$Z(t, x, \omega) := \phi(t, x + Y(\omega), \omega) - Y(\theta(t, \omega)), \quad t \in \mathbf{R}, x \in \mathbf{R}^d, \omega \in \Omega$$

to construct the stable/unstable manifolds. Based on difficult computations using implicit function theorem, perfection arguments and local perturbation of discrete cocycle under the norm

$$\|D\phi\|_\omega := \sup_{n \geq 0} \|D_2\phi(1, Y(\theta(n-1, \omega)), \theta(n-1, \omega))\| e^{n\eta}$$

for small $\eta > 0$.

- Use the continuous-time integrability estimates and the perfect subadditive ergodic theorem to interpolate between discrete time units (or delay periods in SFDE case). This gives asymptotic invariance of the stable/unstable manifolds.

Linearization and Substitution

Assume regularity conditions on the coefficients h, g_i . By the *Substitution Rule*, $\phi(t, Y(\omega), \omega)$ is a stationary *solution* of the *anticipating* Stratonovich SODE

$$\left. \begin{aligned} d\phi(t, Y) &= h(\phi(t, Y)) dt + \sum_{i=1}^m g_i(\phi(t, Y)) \circ dW_i(t), \quad t > 0 \\ \phi(0, Y) &= Y. \end{aligned} \right\} \quad (II)$$

([N-P]).

Linearize the SODE (I) along the stationary trajectory. By substitution, match the solution of the linearized equation with the linearized cocycle $D_2\phi(t, Y(\omega), \omega)$. Hence $D_2\phi(t, Y(\omega), \omega)$, $t \geq 0$, solves the SODE:

$$\left. \begin{aligned} dD_2\phi(t, Y) &= Dh(\phi(t, Y))D_2\phi(t, Y) dt \\ &\quad + \sum_{i=1}^m Dg_i(\phi(t, Y))D_2\phi(t, Y) \circ dW_i(t), \quad t > 0 \\ D_2\phi(0, Y) &= I. \end{aligned} \right\} \quad (III)$$

D_2, D denotes spatial (Fréchet) derivatives.

Similarly, the backward trajectories

$$\phi(t, Y), D_2\phi(t, Y), \quad t < 0,$$

solve the corresponding backward Stratonovich SODE's:

$$\left. \begin{aligned} d\phi(t, Y) &= -h(\phi(t, Y)) dt - \sum_{i=1}^m g_i(\phi(t, Y)) \circ \hat{d}W_i(t), \quad t < 0 \\ \phi(0, Y) &= Y. \end{aligned} \right\} \quad (II^-)$$

$$\left. \begin{aligned} dD_2\phi(t, Y) &= -Dh(\phi(t, Y))D_2\phi(t, Y) dt \\ &\quad - \sum_{i=1}^m Dg_i(\phi(t, Y))D_2\phi(t, Y) \circ \hat{d}W_i(t), \quad t < 0 \\ D_2\phi(0, Y) &= I. \end{aligned} \right\} \quad (III^-)$$

Above SODE's (II)-(III)⁻ give dynamic characterizations of the stable/unstable manifolds.

The following lemma is used to construct the shift-invariant sure event appearing in the statement of the local stable manifold theorem. Gives “perfect versions” of the ergodic theorem and Kingman’s subadditive ergodic theorem.

Lemma 1

(i) Let $h : \Omega \rightarrow \mathbf{R}^+$ be \mathcal{F} -measurable and

$$\int_{\Omega} \sup_{0 \leq u \leq 1} h(\theta(u, \omega)) dP(\omega) < \infty.$$

Then there is a sure event $\Omega_1 \in \mathcal{F}$ such that $\theta(t, \cdot)(\Omega_1) = \Omega_1$ for all $t \in \mathbf{R}$, and

$$\lim_{t \rightarrow \infty} \frac{1}{t} h(\theta(t, \omega)) = 0$$

for all $\omega \in \Omega_1$.

(ii) Suppose $f : \mathbf{R}^+ \times \Omega \rightarrow \mathbf{R} \cup \{-\infty\}$ is a measurable process on (Ω, \mathcal{F}, P) satisfying the following conditions

$$(a) \quad E \sup_{0 \leq u \leq 1} f^+(u) < \infty, \quad E \sup_{0 \leq u \leq 1} f^+(1 - u, \theta(u)) < \infty$$

(b) $f(t_1 + t_2, \omega) \leq f(t_1, \omega) + f(t_2, \theta(t_1, \omega))$ for all $t_1, t_2 \geq 0$ and all $\omega \in \Omega$.

Then there is sure event $\Omega_2 \in \mathcal{F}$ such that $\theta(t, \cdot)(\Omega_2) = \Omega_2$ for all $t \in \mathbf{R}$, and a fixed number $f^* \in \mathbf{R} \cup \{-\infty\}$ such that

$$\lim_{t \rightarrow \infty} \frac{1}{t} f(t, \omega) = f^*$$

for all $\omega \in \Omega_2$.

Proof

[Mo.1], Lemma 7. \square

Theorem 2 ([O], 1968)

Let (Ω, \mathcal{F}, P) be a probability space and $\theta : \mathbf{R}^+ \times \Omega \rightarrow \Omega$ a measurable family of ergodic P -preserving transformations. Let $T : \mathbf{R}^+ \times \Omega \rightarrow L(\mathbf{R}^d)$ be measurable, such that (T, θ) is an $L(\mathbf{R}^d)$ -valued cocycle. Suppose that

$$E \sup_{0 \leq t \leq 1} \log^+ \|T(t, \cdot)\| < \infty, \quad E \sup_{0 \leq t \leq 1} \log^+ \|T(1-t, \theta(t, \cdot))\| < \infty.$$

Then there is a set $\Omega_0 \in \mathcal{F}$ of full P -measure such that $\theta(t, \cdot)(\Omega_0) \subseteq \Omega_0$ for all $t \in \mathbf{R}^+$, and for each $\omega \in \Omega_0$, the limit

$$\lim_{n \rightarrow \infty} [T(t, \omega)^* \circ T(t, \omega)]^{1/(2t)} := \Lambda(\omega)$$

exists in the uniform operator norm. Each $\Lambda(\omega)$ has a non-random spectrum

$$e^{\lambda_1} > e^{\lambda_2} > e^{\lambda_3} > \dots > e^{\lambda_p}$$

where the λ_i 's are distinct. Each e^{λ_i} has a fixed non-random multiplicity m_i and eigen-space $F_i(\omega)$, with $m_i := \dim F_i(\omega)$. Define

$$E_1(\omega) := \mathbf{R}^d, \quad E_i(\omega) := [\oplus_{j=1}^{i-1} F_j(\omega)]^\perp, \quad 1 < i \leq p.$$

Then

$$E_p(\omega) \subset \dots \subset E_{i+1}(\omega) \subset E_i(\omega) \subset \dots \subset E_2(\omega) \subset E_1(\omega) = \mathbf{R}^d,$$

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log \|T(t, \omega)x\| = \lambda_i(\omega) \quad \text{if } x \in E_i(\omega) \setminus E_{i+1}(\omega),$$

and


$$T(t, \omega)(E_i(\omega)) \subseteq E_i(\theta(t, \omega))$$

for all $t \geq 0$, $1 \leq i \leq p$.

Proof.

Based on the discrete version of Oseledec's multiplicative ergodic theorem and Lemma 1. ([Ru.1], I.H.E.S Publications, 1979, pp. 303-304; cf. Furstenberg & Kesten (1960), [Mo.1]), “perfect” infinite-dimensional version and application to regular linear SFDE's. \square

Spectral Theorem




Apply Theorem 2 with $T(t, \omega) := D_2\phi(t, Y(\omega), \omega)$. Then linearized cocycle has random invariant stable and unstable subspaces $\{S(\omega), U(\omega) : \omega \in \Omega\}$:

$$D_2\phi(t, Y(\omega), \omega)(S(\omega)) = S(\theta(t, \omega)),$$

$$D_2\phi(-t, Y(\omega), \omega)(U(\omega)) = U(\theta(-t, \omega)), \quad t \geq 0.$$

[Mo.1].



Estimates on the non-linear cocycle

Theorem 3 ([M-S.2])

There exists a jointly measurable modification of the trajectory random field of (I), denoted by $\{\phi_{s,t}(x) : -\infty < s, t < \infty, x \in \mathbf{R}^d\}$, with the following properties:

Define $\phi : \mathbf{R} \times \mathbf{R}^d \times \Omega \rightarrow \mathbf{R}^d$ by

$$\phi(t, x, \omega) := \phi_{0,t}(x, \omega), \quad x \in \mathbf{R}^d, \omega \in \Omega, t \in \mathbf{R}.$$

Then the following is true for all $\omega \in \Omega$, $\epsilon \in (0, \delta)$:

- (i) For each $x \in \mathbf{R}^d$, and $s, t \in \mathbf{R}$, $\phi_{s,t}(x, \omega) = \phi(t - s, x, \theta(s, \omega))$.
- (ii) (ϕ, θ) is a perfect cocycle:

$$\phi(t + s, \cdot, \omega) = \phi(t, \cdot, \theta(s, \omega)) \circ \phi(s, \cdot, \omega),$$

for all $s, t \in \mathbf{R}$.

- (iii) For each $t \in \mathbf{R}$, $\phi(t, \cdot, \omega) : \mathbf{R}^d \rightarrow \mathbf{R}^d$ is a $C^{k,\epsilon}$ diffeomorphism.
- (iv) The mapping $\mathbf{R}^2 \ni (s, t) \mapsto \phi_{s,t}(\cdot, \omega) \in \text{Diff}^k(\mathbf{R}^d)$ is continuous, where $\text{Diff}^k(\mathbf{R}^d)$ denotes the group of all C^k diffeomorphisms of \mathbf{R}^d , given the C^k -topology.

(v) *The quantities*

$$\begin{aligned}
& \sup_{\substack{0 \leq s, t \leq T, \\ x \in \mathbf{R}^d}} \frac{|\phi_{s,t}(x, \omega)|}{[1 + |x|(\log^+ |x|)^\gamma]}, \quad \sup_{\substack{0 \leq s, t \leq T, \\ x \in \mathbf{R}^d}} \frac{|D_x^\alpha \phi_{s,t}(x, \omega)|}{(1 + |x|^\gamma)}, \\
& \sup_{x \in \mathbf{R}^d} \sup_{\substack{0 \leq s, t \leq T, \\ 0 < |x' - x| \leq \rho}} \frac{|D_x^\alpha \phi_{s,t}(x, \omega) - D_x^\alpha \phi_{s,t}(x', \omega)|}{|x - x'|^\epsilon (1 + |x|)^\gamma}, \\
& \gamma, \rho, T > 0, 1 \leq |\alpha| \leq k
\end{aligned}$$

are finite. The random variables defined by the above expressions have q -th moments for all $q \geq 1$.

$\|\cdot\|_{k,\epsilon} := C^{k,\epsilon}$ -norm on $C^{k,\epsilon}$ mappings $\bar{B}(0,\rho) \rightarrow \mathbf{R}^d$.

Lemma 2

Assume that $\log^+ |Y(\cdot)|$ is integrable. Then

$$\int_{\Omega} \log^+ \sup_{-T \leq t_1, t_2 \leq T} \|\phi(t_2, Y(\theta(t_1, \omega)) + (\cdot), \theta(t_1, \omega))\|_{k,\epsilon} dP(\omega) < \infty \quad (10)$$

for any fixed $0 < T, \rho < \infty$ and any $\epsilon \in (0, \delta)$. The linearized flow $(D_2\phi(t, Y(\omega), \omega), \theta(t, \omega))$, $t \geq 0$, is an $L(\mathbf{R}^d)$ -valued perfect cocycle and

$$\int_{\Omega} \log^+ \sup_{-T \leq t_1, t_2 \leq T} \|D_2\phi(t_2, Y(\theta(t_1, \omega)), \theta(t_1, \omega))\|_{L(\mathbf{R}^d)} dP(\omega) < \infty \quad (11)$$

for any fixed $0 < T < \infty$. The forward cocycle

$(D_2\phi(t, Y(\omega), \omega), \theta(t, \omega), t > 0)$ has a non-random finite Lyapunov spectrum $\{\lambda_p < \dots < \lambda_{i+1} < \lambda_i < \dots < \lambda_2 < \lambda_1\}$. Each Lyapunov exponent λ_i has a non-random multiplicity q_i , $1 \leq i \leq p$, and $\sum_{i=1}^p q_i = d$. The backward linearized cocycle $(D_2\phi(t, Y(\omega), \omega), \theta(t, \omega), t < 0)$, admits a “backward” non-random finite Lyapunov spectrum:

$$\lim_{t \rightarrow -\infty} \frac{1}{t} \log |D_2\phi(t, Y(\omega), \omega)(v(\omega))|, \quad v \in L^0(\Omega, \mathbf{R}^d),$$

taking values $\{-\lambda_i\}_{i=1}^p$ with non-random multiplicities q_i , $1 \leq i \leq p$, and $\sum_{i=1}^p q_i = d$.

The Auxiliary Cocycle

To apply Ruelle's discrete non-linear ergodic theorem ([Ru.1], Theorem 5.1, p. 292), introduce the following auxiliary cocycle $Z : \mathbf{R} \times \mathbf{R}^d \times \Omega \rightarrow \mathbf{R}^d$. This a “centering” of the flow ϕ about the stationary solution:

$$Z(t, x, \omega) := \phi(t, x + Y(\omega), \omega) - Y(\theta(t, \omega))$$

for $t \in \mathbf{R}, x \in \mathbf{R}^d, \omega \in \Omega$.

Lemma 3

(Z, θ) is a perfect cocycle on \mathbf{R}^d and $Z(t, 0, \omega) = 0$ for all $t \in \mathbf{R}$, and all $\omega \in \Omega$.

Proof.

$$\begin{aligned} & Z(t_2, Z(t_1, x, \omega), \theta(t_1, \omega)) \\ &= \phi(t_2, Z(t_1, x, \omega) + Y(\theta(t_1, \omega)), \theta(t_1, \omega)) - Y(\theta(t_2, \theta(t_1, \omega))) \\ &= \phi(t_2, \phi(t_1, x + Y(\omega), \omega), \theta(t_1, \omega)) - Y(\theta(t_2 + t_1, \omega)) \\ &= Z(t_1 + t_2, x, \omega), \quad t_1, t_2 \in \mathbf{R}, \omega \in \Omega, x \in \mathbf{R}^d. \end{aligned}$$

$Z(t, 0, \omega) \equiv 0$ by definition of Z and stationary solution. \square

The proof of the local stable-manifold theorem (Theorem 1) uses a discretization argument that requires the following lemma.

Lemma 4

Suppose that $\log^+ |Y(\cdot)|$ is integrable. Then there is a sure event $\Omega_3 \in \mathcal{F}$ with the following properties:

- (i) $\theta(t, \cdot)(\Omega_3) = \Omega_3$ for all $t \in \mathbf{R}$,
- (ii) For every $\omega \in \Omega_3$ and any $x \in \mathbf{R}^d$, the statement

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log |Z(n, x, \omega)| < 0 \quad (17)$$

implies

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log |Z(t, x, \omega)| = \limsup_{n \rightarrow \infty} \frac{1}{n} \log |Z(n, x, \omega)|. \quad (18)$$

Ruelle's Non-linear Ergodic Theorem

Theorem 4 ([Ru.1], 1979)

Let $\Omega \ni \omega \mapsto F_\omega \in C^{k,\delta}(\mathbf{R}^d, 0; \mathbf{R}^d, 0)$ be measurable such that $E \log^+ \|F_\omega\| \bar{B}(0, 1) < \infty$. Set $F^n(\omega) := F_{\theta(n-1, \omega)} \circ \cdots \circ F_{\theta(1, \omega)} \circ F_\omega$. Suppose $\lambda < 0$ is not in the spectrum of the cocycle $(DF_\omega^n(0), \theta(n, \omega))$. Then there is a sure event $\Omega_0 \in \mathcal{F}$ such that $\theta(1, \cdot)(\Omega_0) \subseteq \Omega_0$, and measurable functions $0 < \alpha(\omega) < \beta(\omega) < 1, \gamma(\omega) > 1$ with the following properties:

(a) If $\omega \in \Omega_0$, the set

$$V_\omega^\lambda := \{x \in \bar{B}(0, \alpha(\omega)) : \|F_\omega^n(x)\| \leq \beta(\omega)e^{n\lambda} \text{ for all } n \geq 0\}$$

is a $C^{k,\delta}$ submanifold of $\bar{B}(0, \alpha(\omega))$.

(b) If $x_1, x_2 \in V_\omega^\lambda$, then

$$\|F_\omega^n(x_1) - F_\omega^n(x_2)\| \leq \gamma(\omega)\|x_1 - x_2\|e^{n\lambda}$$

for all integers $n \geq 0$. If $\lambda' < \lambda$ and $[\lambda', \lambda]$ is disjoint from the spectrum of $(DF_\omega^n(0), \theta(n, \omega))$, then there exists a measurable $\gamma'(\omega) > 1$ such that

$$\|F_\omega^n(x_1) - F_\omega^n(x_2)\| \leq \gamma'(\omega)\|x_1 - x_2\|e^{n\lambda'}$$

for all $x_1, x_2 \in V_\omega^\lambda$ and all integers $n \geq 0$.

Proof

[Ru.1], Theorem 5.1, p. 292.

Construction of the Stable/Unstable Manifolds

Assume the hypotheses of Theorem 1.

Consider the auxiliary cocycle (Z, θ) . Define the family of maps $F_\omega : \mathbf{R}^d \rightarrow \mathbf{R}^d$ by $F_\omega(x) := Z(1, x, \omega)$ for all $\omega \in \Omega$ and $x \in \mathbf{R}^d$. Let $\tau := \theta(1, \cdot) : \Omega \rightarrow \Omega$. Define $F_\omega^n := F_{\tau^{n-1}(\omega)} \circ \cdots \circ F_{\tau(\omega)} \circ F_\omega$. Then cocycle property for Z gives $F_\omega^n = Z(n, \cdot, \omega)$ for each $n \geq 1$. F_ω is $C^{k, \epsilon}$ ($\epsilon \in (0, \delta)$) and $(DF_\omega)(0) = D_2\phi(1, Y(\omega), \omega)$. By measurability of the flow ϕ , the map $\omega \mapsto (DF_\omega)(0)$ is \mathcal{F} -measurable. By (11) of Lemma 2, the map $\omega \mapsto \log^+ \|D_2\phi(1, Y(\omega), \omega)\|_{L(\mathbf{R}^d)}$ is integrable. The discrete cocycle $((DF_\omega^n)(0), \theta(n, \omega), n \geq 0)$ has a non-random Lyapunov spectrum which coincides with that of the linearized continuous cocycle $(D_2\phi(t, Y(\omega), \omega), \theta(t, \omega), t \geq 0)$, viz. $\{\lambda_p < \cdots < \lambda_{i+1} < \lambda_i < \cdots < \lambda_2 < \lambda_1\}$, where each λ_i has fixed multiplicity q_i , $1 \leq i \leq p$ (Lemma 2). If $\lambda_i > 0$ for all $1 \leq i \leq m$, then take $\tilde{\mathcal{S}}(\omega) := \{Y(\omega)\}$ for all $\omega \in \Omega$. Theorem is trivial in this case. Suppose that at least one $\lambda_i < 0$.

Use discrete non-linear ergodic theorem of Ruelle (Theorem 4) and its proof to obtain a sure event $\Omega_1^* \in \mathcal{F}$ such that $\theta(t, \cdot)(\Omega_1^*) = \Omega_1^*$ for all $t \in \mathbf{R}$, \mathcal{F} -measurable positive random variables $\rho_1, \beta_1 : \Omega_1^* \rightarrow (0, 1)$, $\rho_1 < \beta_1$, and a random family of $C^{k, \epsilon}$ ($\epsilon \in (0, \delta)$) submanifolds of $\bar{B}(0, \rho_1(\omega))$ denoted

by $\tilde{\mathcal{S}}_d(\omega)$, $\omega \in \Omega_1^*$, and satisfying the following properties for each $\omega \in \Omega_1^*$:

$$\tilde{\mathcal{S}}_d(\omega) = \{x \in \bar{B}(0, \rho_1(\omega)) : |Z(n, x, \omega)| \leq \beta_1(\omega)e^{(\lambda_{i_0} + \epsilon_1)n} \text{ for all } n \in \mathbf{Z}^+\}. \quad (21)$$

$\tilde{\mathcal{S}}_d(\omega)$ is tangent at 0 to the stable subspace $\mathcal{S}(\omega)$ of the linearized flow $D_2\phi$, viz. $T_0\tilde{\mathcal{S}}_d(\omega) = \mathcal{S}(\omega)$. Therefore $\dim \tilde{\mathcal{S}}_d(\omega)$ is non-random by ergodicity of θ . Also

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log \left[\sup_{\substack{x_1 \neq x_2, \\ x_1, x_2 \in \tilde{\mathcal{S}}_d(\omega)}} \frac{|Z(n, x_1, \omega) - Z(n, x_2, \omega)|}{|x_1 - x_2|} \right] \leq \lambda_{i_0}. \quad (22)$$

The $\theta(t, \cdot)$ -invariant sure event $\Omega_1^* \in \mathcal{F}$ is constructed using the ideas in Ruelle's proof (of Theorem 5.1 in [Ru.1], p. 293), combined with the estimate (10) of Lemma 2 and the subadditive ergodic theorem (Lemma 1 (ii)).

For each $\omega \in \Omega_1^*$, let $\tilde{\mathcal{S}}(\omega)$ be the set defined in part (a) of the theorem. Then by definition of $\tilde{\mathcal{S}}_d(\omega)$ and Z :

$$\tilde{\mathcal{S}}(\omega) = \tilde{\mathcal{S}}_d(\omega) + Y(\omega). \quad (23)$$

Since $\tilde{\mathcal{S}}_d(\omega)$ is a $C^{k, \epsilon}$ ($\epsilon \in (0, \delta)$) submanifold of $\bar{B}(0, \rho_1(\omega))$, then $\tilde{\mathcal{S}}(\omega)$ is a $C^{k, \epsilon}$ ($\epsilon \in (0, \delta)$) submanifold of $\bar{B}(Y(\omega), \rho_1(\omega))$. Furthermore, $T_{Y(\omega)}\tilde{\mathcal{S}}(\omega) = T_0\tilde{\mathcal{S}}_d(\omega) = \mathcal{S}(\omega)$. Hence $\dim \tilde{\mathcal{S}}(\omega) = \dim \mathcal{S}(\omega) = \sum_{i=i_0}^p q_i$, and is non-random.

Now (22) implies that

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log |Z(n, x, \omega)| \leq \lambda_{i_0} \quad (24)$$

for all ω in Ω_1^* and all $x \in \tilde{\mathcal{S}}_d(\omega)$. Therefore by Lemma 4, there is a sure event $\Omega_2^* \subseteq \Omega_1^*$ such that $\theta(t, \cdot)(\Omega_2^*) = \Omega_2^*$ for all $t \in \mathbf{R}$, and

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log |Z(t, x, \omega)| \leq \lambda_{i_0} \quad (25)$$

for all $\omega \in \Omega_2^*$ and all $x \in \tilde{\mathcal{S}}_d(\omega)$. Therefore (2) holds.

To prove (b), let $\omega \in \Omega_1^*$. By (22), there is a positive integer $N_0 := N_0(\omega)$ (independent of $x \in \tilde{\mathcal{S}}_d(\omega)$) such that $Z(n, x, \omega) \in \bar{B}(0, 1)$ for all $n \geq N_0$. Let $\Omega_4^* := \Omega_2^* \cap \Omega_3$, where Ω_3 is the shift-invariant sure event defined in the proof of Lemma 4. Then Ω_4^* is a sure event and $\theta(t, \cdot)(\Omega_4^*) = \Omega_4^*$ for all $t \in \mathbf{R}$. By cocycle property, Mean-Value theorem and the ergodic theorem (Lemma 1(i)), we get (b).

To prove the invariance property (4), apply the Osledec theorem to the cocycle $(D_2\phi(t, Y(\omega), \omega), \theta(t, \omega))$ ([Mo.1], Theorem 4, Corollary 2). This gives a sure $\theta(t, \cdot)$ -invariant event, also denoted by Ω_1^* , such that

$D_2\phi(t, Y(\omega), \omega)(\mathcal{S}(\omega)) \subseteq \mathcal{S}(\theta(t, \omega))$ for all $t \geq 0$ and all $\omega \in \Omega_1^*$. Equality holds because $D_2\phi(t, Y(\omega), \omega)$ is injective and $\dim \mathcal{S}(\omega) = \dim \mathcal{S}(\theta(t, \omega))$ for all $t \geq 0$ and all $\omega \in \Omega_1^*$.

To prove the asymptotic invariance property (3), use the ideas in the proofs of Theorems 5.1 and 4.1 in [Ru.1], pp. 285-297, to pick random variables ρ_1, β_1 and a sure event (also denoted by) Ω_1^* such that $\theta(t, \cdot)(\Omega_1^*) = \Omega_1^*$ for all $t \in \mathbf{R}$, and

$$\rho_1(\theta(t, \omega)) \geq \rho_1(\omega)e^{(\lambda_{i_0} + \epsilon_1)t}, \quad \beta_1(\theta(t, \omega)) \geq \beta_1(\omega)e^{(\lambda_{i_0} + \epsilon_1)t} \quad (26)$$

for all $t \geq 0, \omega \in \Omega_1^*$. Use (b) to obtain a sure event $\Omega_5^* \subseteq \Omega_4^*$ such that $\theta(t, \cdot)(\Omega_5^*) = \Omega_5^*$ for all $t \in \mathbf{R}$, and for any $0 < \epsilon < \epsilon_1$ and $\omega \in \Omega_4^*$, there exists $\beta^\epsilon(\omega) > 0$ (independent of x) with

$$|\phi(t, x, \omega) - Y(\theta(t, \omega))| \leq \beta^\epsilon(\omega)e^{(\lambda_{i_0} + \epsilon)t} \quad (27)$$

for all $x \in \tilde{\mathcal{S}}(\omega)$, $t \geq 0$. Fix $t \geq 0$, $\omega \in \Omega_5^*$ and $x \in \tilde{\mathcal{S}}(\omega)$. Let n be a non-negative integer. Then the cocycle property and (27) imply that

$$\begin{aligned} & |\phi(n, \phi(t, x, \omega), \theta(t, \omega)) - Y(\theta(n, \theta(t, \omega)))| \\ &= |\phi(n + t, x, \omega) - Y(\theta(n + t, \omega))| \\ &\leq \beta^\epsilon(\omega)e^{(\lambda_{i_0} + \epsilon)(n+t)} \\ &\leq \beta^\epsilon(\omega)e^{(\lambda_{i_0} + \epsilon)t}e^{(\lambda_{i_0} + \epsilon_1)n}. \end{aligned} \quad (28)$$

If $\omega \in \Omega_5^*$, then it follows from (26), (27), (28) and the definition of $\tilde{\mathcal{S}}(\theta(t, \omega))$ that there exists $\tau_1(\omega) > 0$ such that

$\phi(t, x, \omega) \in \tilde{\mathcal{S}}(\theta(t, \omega))$ for all $t \geq \tau_1(\omega)$. This proves asymptotic invariance.

Prove (d) by running both the flow ϕ and the shift θ backward in time:

$$\tilde{\phi}(t, x, \omega) := \phi(-t, x, \omega), \quad \tilde{Z}(t, x, \omega) := Z(-t, x, \omega), \quad \tilde{\theta}(t, \omega) := \theta(-t, \omega)$$

for all $t \geq 0$ and all $\omega \in \Omega$. $(\tilde{Z}(t, \cdot, \omega), \tilde{\theta}(t, \omega), t \geq 0)$ is a smooth cocycle, with $\tilde{Z}(t, 0, \omega) = 0$ for all $t \geq 0$. The linearized flow $(D_2\tilde{\phi}(t, Y(\omega), \omega), \tilde{\theta}(t, \omega), t \geq 0)$ is an $L(\mathbf{R}^d)$ -valued perfect cocycle with a non-random finite Lyapunov spectrum $\{-\lambda_1 < -\lambda_2 < \dots < -\lambda_i < -\lambda_{i+1} < \dots < -\lambda_p\}$ where $\{\lambda_p < \dots < \lambda_{i+1} < \lambda_i < \dots < \lambda_2 < \lambda_1\}$ is the Lyapunov spectrum of the forward linearized flow $(D_2\phi(t, Y(\omega), \omega), \theta(t, \omega), t \geq 0)$. Apply first part of the proof to get *stable manifolds* for the backward flow $\tilde{\phi}$ satisfying assertions (a), (b), (c). This translates into the existence of *unstable manifolds* for the original flow ϕ , and (d), (e), (f) automatically hold. Hence there is a sure event $\Omega_6^* \in \mathcal{F}$ such that $\theta(-t, \cdot)(\Omega_6^*) = \Omega_6^*$ for all $t \in \mathbf{R}$, and (d), (e) and (f) hold for all $\omega \in \Omega_6^*$.

Define the sure event $\Omega^* := \Omega_6^* \cap \Omega_5^*$. Then $\theta(t, \cdot)(\Omega^*) = \Omega^*$ for all $t \in \mathbf{R}$. Assertions (a)-(f) hold for all $\omega \in \Omega^*$.

Measurability of the stable manifolds follows from

$$\tilde{\mathcal{S}}(\omega) = Y(\omega) + \tilde{\mathcal{S}}_d(\omega) \tag{29}$$

$$\tilde{\mathcal{S}}_d(\omega) = \lim_{m \rightarrow \infty} \bar{B}(0, \rho_1(\omega)) \cap \bigcap_{i=1}^m f_i(\cdot, \omega)^{-1}(\bar{B}(0, 1)) \quad (30)$$

$$f_n(x, \omega) := \beta_1(\omega)^{-1} e^{-(\lambda_{i_0} + \epsilon_1)n} Z(n, x, \omega), \quad x \in \mathbf{R}^d, \omega \in \Omega_1^*,$$

for all integers $n \geq 0$. (Above limit is taken in the metric d^* on $\mathcal{C}(\mathbf{R}^d)$.) Use joint continuity of translation and measurability of Y , f_i , ρ_1 , finite intersections and the continuity of the maps

$$\mathbf{R}^+ \ni r \mapsto \bar{B}(0, r) \in \mathcal{C}(\mathbf{R}^d).$$

$$\text{Hom}(\mathbf{R}^d) \ni f \mapsto f^{-1}(\bar{B}(0, 1)) \in \mathcal{C}(\mathbf{R}^d).$$

When h, g_i are in C_b^∞ , adapt above argument to give a sure event in \mathcal{F} , also denoted by Ω^* such that $\tilde{\mathcal{S}}(\omega), \tilde{\mathcal{U}}(\omega)$ are C^∞ for all $\omega \in \Omega^*$. \square

Remarks

- (i) Replace the stationary random variable Y by its invariant distribution ρ , with $\int_{\mathbf{R}^d} |x| d\rho(x) < \infty$. Formulate result with respect to the product measure $P \otimes \rho$ and the underlying skew-product flow. This would give stable and unstable manifolds that are defined a.e. $(P \otimes \rho)$; cf. [C] for the globally asymptotically stable case on a compact manifold.
- (ii) In Stratonovich SODE (I), replace global boundedness on g'_i s by requiring

$$\mathbf{R}^d \ni x \mapsto \sum_{l=1}^m \frac{\partial g_l^i(x)}{\partial x_j} g_l^j(x) \in \mathbf{R}, \quad 1 \leq i, j \leq d$$

to be in $C_b^{k,\delta}$.

- (iii) Conjecture: The global boundedness condition is not needed. This conjecture is not hard to check if the vector fields g_i , $1 \leq i \leq m$, are C_b^∞ and generate a finite-dimensional solvable Lie algebra. See [Ku], Theorem 4.9.10, p. 212.
- (iv) Theorem holds for the Itô SODE

$$dx(t) = h(x(t)) dt + \sum_{i=1}^m g_i(x(t)) dW_i(t), \quad (II)$$

with $h, g_i : \mathbf{R}^d \rightarrow \mathbf{R}^d, 1 \leq i \leq m$, in $C_b^{k,\delta}$.

(v) Replace (I) with Kunita-type SODE

$$\left. \begin{aligned} d\phi(t) &= \overset{\circ}{F}(\circ dt, \phi(t)), \quad t > s \\ \phi(s) &= x \end{aligned} \right\}$$

where $\overset{\circ}{F}$ is a spatial semimartingale helix (i.e. with stationary ergodic increments) and with local characteristics of class $(B_{ub}^{k+1,\delta}, B_{ub}^{k,\delta})$ and the function

$$[0, \infty) \times \mathbf{R}^d \ni (t, x) \mapsto \sum_{j=1}^d \frac{\partial a^{\cdot,j}(t, x, y)}{\partial x_j} \Big|_{y=x} \in \mathbf{R}^d$$

belongs to $B_{ub}^{k,\delta}$. In the Itô case, last condition is not needed.

$$\begin{aligned} \overset{\circ}{F}(t, x) &= V(t, x) + M(t, x) \\ a^{i,j}(t, x, y) &:= \frac{d}{dt} \langle M^i(\cdot, x), M^j(\cdot, y) \rangle(t) \\ b^i(t, x) &:= \frac{d}{dt} V^i(t, x), \quad x, y \in \mathbf{R}^d, 1 \leq i, j \leq d \end{aligned}$$

Proof of Theorem 3

Cocycle property (ii): approximate the flow using helix mollifiers of Brownian motion:

$$W^k(t) := k \int_{t-1/k}^t W(s) ds - k \int_{-1/k}^0 W(s) ds.$$

$$W^k(t_2, \theta(t_1, \omega)) = W^k(t_1 + t_2, \omega) - W^k(t_1, \omega), \quad k \geq 1$$

([I-W], cf. [Mo.1], [Mo.2] for linear infinite-dimensional case).

(iii) and (iv) are well-known to hold for a.a. $\omega \in \Omega$ ([Ku], Theorem 4.6.5).

A perfect version of $\phi_{s,t}$ satisfying (i)-(iv) for *all* $\omega \in \Omega$, is obtained in [A-S] by perfection techniques and the diffeomorphism theorem for flows ([Ku], Theorem 4.6.5; cf. also [M-S.1]).

By [M-S.2], the random variables

$$X_1 := \sup_{\substack{0 \leq s \leq t \leq T, \\ x \in \mathbf{R}^d}} \frac{|\phi_{s,t}(x, \cdot)|}{[1 + |x|(\log^+ |x|)^\gamma]},$$

$$X_2 := \sup_{\substack{0 \leq s \leq t \leq T, \\ x \in \mathbf{R}^d}} \frac{|x|}{[1 + |\phi_{s,t}(x, \cdot)|(\log^+ |x|)^\gamma]}$$

have q -th moments for all $q \geq 1$. It is sufficient to show that the random variable

$$\hat{X}_1 := \sup_{\substack{0 \leq s \leq t \leq T, \\ x \in \mathbf{R}^d}} \frac{|\phi_{t,s}(x, \cdot)|}{[1 + |x|(\log^+ |x|)^\gamma]}$$

has q -th moments for all $q \geq 1$. Assume (without loss of generality) that $\gamma \in (0, 1)$. From the definition of X_2 ,

$$|y| \leq X_2[1 + |\phi_{s,t}(y, \cdot)|(\log^+ |y|)^\gamma]$$

for all $0 \leq s \leq t \leq T, y \in \mathbf{R}^d$. Use the substitution

$$y = \phi_{t,s}(x, \omega) = \phi_{s,t}^{-1}(x, \omega), \quad \phi_{s,t}(y, \omega) = x, \quad 0 \leq s \leq t \leq T, \omega \in \Omega, x \in \mathbf{R}^d,$$

to rewrite above inequality as

$$|y| \leq X_2[1 + |x|(\log^+ |y|)^\gamma].$$

Solve above inequality (by taking \log^+) for $\log^+ |y|$. Therefore, there exists a non-random constant $K_1 := K_1(\gamma) > 0$ such that

$$|y| \leq K_1 X_2[1 + |x|\{1 + (\log^+ |X_2|)^\gamma + (\log^+ |x|)^\gamma\}].$$

Since X_2 has moments of all orders, the above inequality implies that \hat{X}_1 also has q -th moments for all $q \geq 1$.

Complete proof by [Ku], [M-S.2] and GRR. \square

Proof of Lemma 2

We first prove (11). Start with the perfect cocycle property for (ϕ, θ) :

$$\phi(t_1 + t_2, \cdot, \omega) = \phi(t_2, \cdot, \theta(t_1, \omega)) \circ \phi(t_1, \cdot, \omega) \quad (12)$$

for all $t_1, t_2 \in \mathbf{R}$ and all $\omega \in \Omega$. Cocycle property for $(D_2\phi(t, Y(\omega), \omega), \theta(t, \omega))$ follows directly by taking Fréchet derivatives at $Y(\omega)$ on both sides of (12); viz.

$$\begin{aligned} D_2\phi(t_1 + t_2, Y(\omega), \omega) \\ &= D_2\phi(t_2, \phi(t_1, Y(\omega), \omega), \theta(t_1, \omega)) \circ D_2\phi(t_1, Y(\omega), \omega) \\ &= D_2\phi(t_2, Y(\theta(t_1, \omega)), \theta(t_1, \omega)) \circ D_2\phi(t_1, Y(\omega), \omega) \end{aligned} \quad (13)$$

for all $\omega \in \Omega, t_1, t_2 \in \mathbf{R}$. Existence of a fixed discrete spectrum for $D_2\phi(t, Y)$ follows from [Mo.1] and [M-S.1], using the integrability property (11) and the ergodicity of θ . ((11) follows from (13) and Theorem 3 (v)). But (10) implies (11)! Therefore it is sufficient to prove (10).

In view of (1) and the identity

$$\phi_{t_1, t_1+t_2}(x, \omega) = \phi(t_2, x, \theta(t_1, \omega)), \quad x \in \mathbf{R}^d, t_1, t_2 \in \mathbf{R},$$

(Theorem 3(i)), (10) (for $\epsilon = 0$) will follow from

$$\int_{\Omega} \log^+ \sup_{\substack{0 \leq s, t \leq T, \\ |x'| \leq \rho}} |D_x^\alpha \phi_{s,t}(\phi_{0,s}(Y(\omega), \omega) + x', \omega)| dP(\omega) < \infty, \quad 0 \leq |\alpha| \leq k. \quad (14)$$

Denote random “constants” by $K_i, i = 1, 2, 3, 4$. Each $K_i := K_i(\rho, T), i = 1, 2, 3, 4$, has q -th moments for all $q \geq 1$. The following inequalities follow easily from Theorem 3 (v).

$$\begin{aligned} & \log^+ \sup_{\substack{s, t \in [0, T], \\ |x'| \leq \rho}} |D_x^\alpha \phi_{s,t}(\phi_{0,s}(Y(\omega), \omega) + x', \omega)| \\ & \leq \log^+ \sup_{s \in [0, T]} \{K_1(\omega)[1 + (\rho + |\phi_{0,s}(Y(\omega), \omega)|)^2]\} \\ & \leq \log^+ K_2(\omega) + \log^+[1 + 2\rho^2 + K_3(\omega)(1 + |Y(\omega)|^4)] \\ & \leq \log^+ K_4(\omega) + \log[1 + 2\rho^2] + 4 \log^+ |Y(\omega)| \end{aligned} \quad (15)$$

for all $\omega \in \Omega$. (15)+ integrability hypothesis on Y imply (14). \square

Proof of Lemma 4

The integrability condition (10) of Lemma 2 implies that

$$\int_{\Omega} \log^+ \sup_{\substack{0 \leq t_1, t_2 \leq 1, \\ x^* \in \bar{B}(0,1)}} \|D_2 Z(t_1, x^*, \theta(t_2, \omega))\|_{L(\mathbf{R}^d)} dP(\omega) < \infty. \quad (19)$$

Therefore by (the perfect version of) the ergodic theorem (Lemma 1(i)), there is a sure event $\Omega_3 \in \mathcal{F}$ such that $\theta(t, \cdot)(\Omega_3) = \Omega_3$ for all $t \in \mathbf{R}$, and

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log^+ \sup_{\substack{0 \leq u \leq 1, \\ x^* \in \bar{B}(0,1)}} \|D_2 Z(u, x^*, \theta(t, \omega))\|_{L(\mathbf{R}^d)} = 0 \quad (20)$$

for all $\omega \in \Omega_3$.

Let $\omega \in \Omega_3$ and suppose $x \in \mathbf{R}^d$ satisfies (17). Then (17) implies that there exists a positive integer $N_0(x, \omega)$ such that $Z(n, x, \omega) \in \bar{B}(0, 1)$ for all $n \geq N_0$. Let $n \leq t < n + 1$, $n \geq N_0$. Then by the cocycle property for (Z, θ) and the Mean Value Theorem:

$$\begin{aligned} & \sup_{n \leq t \leq n+1} \frac{1}{t} \log |Z(t, x, \omega)| \\ & \leq \frac{1}{n} \log^+ \sup_{\substack{0 \leq u \leq 1, \\ x^* \in \bar{B}(0,1)}} \|D_2 Z(u, x^*, \theta(n, \omega))\|_{L(\mathbf{R}^d)} + \frac{n}{(n+1)} \frac{1}{n} \log |Z(n, x, \omega)|. \end{aligned}$$

Take $\limsup_{n \rightarrow \infty}$ in the above relation and use (20) to get

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log |Z(t, x, \omega)| \leq \limsup_{n \rightarrow \infty} \frac{1}{n} \log |Z(n, x, \omega)|.$$

The inequality

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log |Z(n, x, \omega)| \leq \limsup_{t \rightarrow \infty} \frac{1}{t} \log |Z(t, x, \omega)|,$$

is obvious. Hence (18) holds. \square

References

- [A-S] Arnold, L., and Scheutzow, M. K. R., Perfect cocycles through stochastic differential equations, *Probab. Th. Rel. Fields*, 101, (1995), 65-88.
- [A-I] Arnold, L., and Imkeller, P., Stratonovich calculus with spatial parameters and anticipative problems in multiplicative ergodic theory, *Stoch. Proc. Appl.* 62 (1996), 19-54.
- [C] Carverhill, A., Flows of stochastic dynamical systems: Ergodic theory, *Stochastics*, 14 (1985), 273-317.
- [I-W] Ikeda, N., and Watanabe, S., *Stochastic Differential Equations and Diffusion Processes*, Second Edition, North-Holland-Kodansha (1989).
- [Ku] Kunita, H., *Stochastic Flows and Stochastic Differential Equations*, Cambridge University Press, Cambridge, New York, Melbourne, Sydney (1990).
- [M-N-S] Millet, A., Nualart, D., and Sanz, M., Large deviations for a class of anticipating stochastic differential equations, *The Annals of Probability*, 20 (1992), 1902-1931.
- [Mo.1] Mohammed, S.-E. A., The Lyapunov spectrum and stable manifolds for stochastic linear delay equations, *Stochastics and Stochastic Reports*, Vol. 29 (1990), 89-131.
- [Mo.2] Mohammed, S.-E. A., Lyapunov exponents and stochastic flows of linear and affine hereditary Systems, *Diffusion Processes and Related Problems in Analysis*, Vol. II,

edited by Mark Pinsky and Volker Wihstutz, Birkhauser (1992), 141-169.

- [M-S.1] Mohammed, S.-E. A., and Scheutzow, M. K. R., Lyapunov exponents of linear stochastic functional differential equations driven by semimartingales, Part I: The multiplicative ergodic theory, *Ann. Inst. Henri Poincaré, Probabilités et Statistiques*, Vol. 32, 1, (1996), 69-105. pp. 43.
- [M-S.2] Mohammed, S.-E. A., and Scheutzow, M. K. R., Spatial estimates for stochastic flows in Euclidean space, *The Annals of Probability*, 26, 1, (1998), 56-77.
- [Nu] Nualart, D., Analysis on Wiener space and anticipating stochastic calculus, *St. Flour Notes*.
- [N-P] Nualart, D., and Pardoux, E., Stochastic calculus with anticipating integrands, Analysis on Wiener space and anticipating stochastic calculus, *Probab. Th. Rel. Fields*, 78 (1988), 535-581.
- [O] Oseledec, V. I., A multiplicative ergodic theorem. Lyapunov characteristic numbers for dynamical systems, *Trudy Moskov. Mat. Obšč.* 19 (1968), 179-210. English transl. *Trans. Moscow Math. Soc.* 19 (1968), 197-221.
- [O-P] Ocone, D., and Pardoux, E., A generalized Itô-Ventzell formula. Application to a class of anticipating stochastic differential equations, *Ann. Inst. Henri Poincaré, Probabilités et Statistiques*, Vol. 25, no. 1 (1989), 39-71.

- [Ru.1] Ruelle, D., Ergodic theory of differentiable dynamical systems, *Publ. Math. Inst. Hautes Etud. Sci.* (1979), 275-306.
- [Ru.2] Ruelle, D., Characteristic exponents and invariant manifolds in Hilbert space, *Annals of Mathematics* 115 (1982), 243–290.
- [Sc] Scheutzow, M. K. R., On the perfection of crude cocycles, *Random and Computational Dynamics*, 4, (1996), 235-255.
- [Wa] Wanner, T., Linearization of random dynamical systems, in *Dynamics Reported*, vol. 4, edited by U. Kirchgraber and H.O. Walther, Springer (1995), 203-269.